

WILDWOOD FAMILY CLINIC, S.C.

"HEALTHCARE FOR ALL AGES"

MADISON OFFICE 4901 Cottage Grove Road Madison, WI 53716 (608) 221-1501

COTTAGE GROVE OFFICE 251 East Cottage Grove Road Cottage Grove, WI 53527 (608) 839-3515

Children's Ibuprofen Dosage Chart

DO NOT GIVE TO PATIENT'S UNDER 6 MONTHS OF AGE

Repeat every 6 – 8 hours as needed or as recommended by your healthcare provider

Weight	Infant Drops	Children's	Children's Junior Strength Caplets	
	(50 mg/1.25	Liquid *	Chewable	or Chewable Tablets
	mL syringe)	100 mg/	Tablets	(100 mg tablets)
		5 mL	(50 mg tablets)	
6 - 11 lbs. / 2.7-5 kg	½ syringe (0.625 mL)			
10 17 lbs / 5 4 7 7 lbs		17.1		
12-17 lbs / 5.4-7.7 kg	1 syringe	½ teaspoon		
	(1.25 mL)	(2.5 mL)		
18-23 lbs / 8-10.5 kg	1 ½ syringe	3/4 teaspoon	1 ½ Tablets	
	(1.875 mL)	(3.75 mL)		
24-35 lbs / 11-16 kg	2 syringes (2	1 teaspoon	2 Tablets	1 Tablet
	x 1.25 = 2.5	(5 mL)		
	mL)	(0 1112)		
36-47 lbs / 16.4-21.4		1 1/2 teaspoons	3 Tablets	1 1/2 Tablets
kg		(7.5 mL)	0 1 4 5 10 10	1 /2 1 401010
48 - 59 lbs / 22-27 kg		2 teaspoons	4 Tablets	2 Tablets
		(10 mL)	4 1 4 5 10 10 10	2 Tablets
60 - 71 lbs / 27.2-32		2 ½ teaspoons	5 Tablets	2 ½ Tablets
kg		(12.5 mL)	o Tabloto	2 /2 1 abicts
72 - 95 lbs / 33-43 kg		3 teaspoons	6 Tablets	3 Tablets
72 00 150 7 00 40 kg			o rabiets	3 Tablets
Over 05 lbs / >42 lcs		(15 mL)	0.11.	
Over 95 lbs / >43 kg	*	4 teaspoons	8 tablets	4 Tablets
		(20 mL)		

95 lbs/43 kg. and over - May use 2 regular strength (200 mg) adult ibuprofen tablets/capsules.

* USE ORAL SYRINGES OR SUPPLIED MEDICINE CUP TO MEASURE LIQUID (100 MG/ 5 ML)
- NOT HOUSEHOLD TEASPOONS WHICH CAN DIFFER IN SIZE
Do not use aspirin in children during viral illnesses because of association with

Reye's syndrome.